NATURAL PROGRESSION

URBAN EDGE environmental CONSULTING

Habitats Regulations Assessment for the Mid Sussex District Plan

Appropriate Assessment Report for the Consultation Draft District Plan

October 2014



NATURAL PROGRESSION



Habitats Regulations Assessment for the Mid Sussex District Plan

Appropriate Assessment Report for the

Consultation Draft District Plan

Client:	Mid Sussex District Council			
Report No.:	UE-0158 Mid Sussex DP HRA_5_141010			
Version:	5			
Status:	Final			
Date:	October 2014			
Author:	NP/SP			
Checked:	SP			
Approved:	NP			

Contents

Exec	utive Summary	i
E1	Introduction	i
E2	Findings of the Screening Stage	i
E3	Conclusions of the Appropriate Assessment	ii
1 1	Introduction	1
1.1	Purpose of this Report	1
1.2	The Mid Sussex District Plan	1
1.3	Habitats Regulations Assessment	2
1.4	Structure of this Document	2
1.5	Relationship to Other Documents	3
2	Methodology	5
2.1	Guidance and Best Practice	5
2.2	Methodology	6
2.3	Screening	6
2.4	Appropriate Assessment	8
2.5	In Combination Effects	8
3 I	European Sites	9
3.1	Scope of the Assessment	9
3.2	Qualifying Features	9
3.3	Conservation Objectives	9
3.4	Condition Status	12
4 I	European Site Characterisation	13
4.1	Qualifying Species of the Special Area of Conservation	13
4.2	Qualifying Habitats of the Special Area of Conservation	13
4.3	SPA Bird Populations	14
5	Atmospheric Pollution Impact Pathways	17
5.1	Introduction	17

5.2	Understanding the Vulnerability of SAC Habitats to Atmospheric Pollution	17
5.3	Evidence for Pollution Impacts within Ashdown Forest	18
5.4	Contribution of the District Plan to Future Pollution Deposition within Ashdown Forest	19
5.5	Other Plans and Projects acting In Combination	23
5.6	Avoidance and Mitigation Measures proposed by the District Plan	24
5.7	Impact Assessment	25
5.8	Conclusions regarding Adverse Effects on Integrity	26
6 Di	sturbance Impact Pathways	27
6.1	Introduction	27
6.2	Understanding the Vulnerability of SPA Species to Disturbance	27
6.3	Evidence for Disturbance Impacts within Ashdown Forest	29
6.4	Determining a Zone of Influence for Visiting Patterns to Ashdown Forest	30
6.5	Contribution of the District Plan to Future Visiting Rates within Ashdown Forest	32
6.6	Other Plans and Projects acting In Combination	35
6.7	Avoidance and Mitigation Measures proposed by the District Plan	35
6.8	Impact Assessment	36
6.9	Conclusions regarding Adverse Effects on Integrity	37
7 Su	Immary and Conclusions	39
7.1	Summary	39
7.2	Conclusions	39
7.3	Consultation Arrangements	39
7.4	Next Steps	39
Refere	nces and Bibliography	41
Appen	dix I: Updated Screening Assessment	A



List of Tables and Figures

Table 2.1: Stages in HRA drawing on guidance from DCLG and Natural England	7
Table 3.1: Qualifying features of the Ashdown Forest SAC and SPA	11
Table 3.2: Conservation objectives for Ashdown Forest SAC and SPA	11
Table 3.3: Typical species of Annex 1 habitat types present within SAC	12
Table 3.4: SSSI condition summary by area (1 August 2014)	12
Table 5.1: Critical load for Ashdown Forest SAC compared to actual maximum load	19
Table 6.1: Predicted additional visitor rates to the SPA per 16 daylight hours in Sept. resu development at different locations (Source: Clarke et al. 2010)	Ilting from new 32
Figure 3.1: Ashdown Forest SAC/SPA in relation to Mid Sussex district	10
Figure 5.1: APIS grid reference locations	20
Figure 5.2: Major roads crossing Ashdown Forest	22
Figure 6.1: Cumulative frequency of car visitors by distance, for the Dorset Heaths, Thame and Ashdown Forest (Source: Clarke et al. 2010)	es Basin Heaths 31
Figure 6.2: Distances travelled to Ashdown Forest in 2008 by all respondents who gave a (n=286)	full post code 33
Figure 6.3: Ashdown Forest's 7km Zone of Influence within Mid Sussex	34



Abbreviations

- AA Appropriate Assessment
- AADT Annual Average Daily Traffic
- HDV Heavy Duty Vehicle
- HRA Habitats Regulations Assessment
- IROPI Imperative Reasons of Overriding Public Interest
- keq/ha/yr Kilograms equivalent per hectare per year
- Kg/ha/yr Kilograms per hectare per year
- LDD Local Development Document
- LNR Local Nature Reserve
- NGR National Grid Reference
- NO_X Oxides of nitrogen
- SAC Special Area of Conservation
- SAMM Strategic Access Management and Monitoring
- SANG Suitable Alternative Natural Greenspace
- SNCI Site of Nature Conservation Importance
- SPA Special Protection Area
- SSSI Site of Special Scientific Interest



Executive Summary

E1 Introduction

- E1.1 This document sets out the Habitats Regulations Assessment for the Consultation Draft District Plan for Mid Sussex. It draws on information previously published in other documents which form part of the overall HRA procedure for the District Plan, including research reports and earlier versions of the HRA.
- E1.2 The outputs of the report include information in relation to:
 - Chapter 2 discusses the methodology for assessment, and updates the findings of the screening stage;
 - Chapter 3 defines relevant European sites, their qualifying features and conservation objectives;
 - Chapter 4 discusses the characteristics of European sites including available information about the population and ecology of qualifying species, and descriptions of qualifying habitats;
 - Chapter 5 presents available evidence regarding the mechanisms of atmospheric pollution impacts, and undertakes an assessment of these impacts in view of the sites' conservation objectives;
 - Chapter 6 presents available evidence regarding the mechanisms of disturbance impacts, and undertakes an assessment of these impacts in view of the sites' conservation objectives;
 - Chapter 7 presents a summary and concludes the document.

E2 Findings of the Screening Stage

- E2.1 The District Plan's forerunner, the draft Core Strategy, underwent an HRA screening and scoping exercise in late 2007 and early 2008. This found that five designated areas were potentially at risk of effects from the Plan:
 - Ashdown Forest Special Area of Conservation (SAC);
 - Ashdown Forest Special Protection Area (SPA);
 - Castle Hill SAC;
 - Lewes Downs SAC; and
 - Mole Gap to Reigate Escarpment SAC.



- E2.2 Castle Hill, Lewes Downs and Mole Gap to Reigate Escarpment were screened out of the assessment, largely due to their distance from the district and the low likelihood of residents travelling along roads close to the sites. Natural England (2008) concurred with these findings in its screening opinion on the Core Strategy. Acknowledging that the plan is not necessary to the management of any European site, the screening exercise found likely significant effects on Ashdown Forest SAC/SPA as a result of disturbance and atmospheric pollution.
- E2.3 The updated screening assessment for the Consultation Draft District Plan found that the following proposed policies were likely to significantly affect Ashdown Forest SAC or SPA. They were taken forward for more detailed Appropriate Assessment.
 - DP2 Sustainable Economic Development;
 - DP3 Town Centre Development;
 - DP5 Housing;
 - > DP7 Strategic Allocation to the east of Burgess Hill at Kings Way;
 - > DP8 Strategic Allocation to the north and north-west of Burgess Hill;
 - DP12 New Homes in the Countryside;
 - DP29 Rural Exception Sites; and
 - > DP30 Gypsies, Travellers and Travelling Showpeople.

E3 Conclusions of the Appropriate Assessment

- E3.1 Ashdown Forest is designated as an SAC for its European dry heaths and North Atlantic wet heaths with *E. tetralix*. Great crested newts are present in qualifying numbers but are not a primary reason for site selection. Predicted traffic growth as a result of development proposed by the District Plan is below the significance threshold set for European sites. As a result, it can be concluded that the District Plan will not result in adverse effects on the ecological integrity of the Ashdown Forest SAC.
- E3.2 Ashdown Forest is classified as an SPA for its breeding populations of Dartford warbler and nightjar, which are listed on Annex 1 of the Birds Directive. Breeding woodlark (also an Annex 1 species) are present in qualifying numbers. The Council is committed to delivering a strategic Suitable Alternative Natural Greenspace (SANG) at Ashplats Wood and East Court, East Grinstead, to avoid increases in recreational pressure resulting from planned residential developments within the District Plan. Additionally, the Council has adopted an interim Strategic Access Management and Monitoring (SAMM) Strategy to mitigate the residual effects of visitors originating from within the district, and is committed to replacing this with a joint SAMM Strategy prepared in collaboration with the Ashdown Forest Conservators, Natural England and other affected authorities. It can be concluded that the District Plan will not result in adverse effects on the ecological integrity of the Ashdown Forest SPA.

1 Introduction

1.1 Purpose of this Report

- 1.1.1 This report has been prepared for Mid Sussex District Council as part of the Habitats Regulations Assessment (HRA) for the District Plan. The report accompanies the Consultation Draft version of the Plan and forms part of the evidence base upon which the Plan is based.
- 1.1.2 HRA is a requirement of the Conservation of Habitats and Species Regulations 2010 (as amended; 'the Habitats Regulations'). The assessment focuses on the likely significant effects of the plan on the nature conservation interests of European-protected areas in and around the district, and seeks to establish whether or not there will be any adverse effects on the ecological integrity of these European sites as a result of proposals in the plan.

1.2 The Mid Sussex District Plan

- 1.2.1 The Mid Sussex District Plan will provide the overall planning policy framework for the district for a period of 17 years between 2014 and 2031. A Housing and Economic Development Needs Assessment will provide the objectively assessed housing need within Mid Sussex over the plan period. The Council's Strategic Housing Land Availability Assessment indicates the overall supply of land that could be available for residential development over the period. A primary role for the District Plan is to reconcile differences between housing need and land supply, taking account of the need to cooperate with neighbouring authorities, in the context of environmental and infrastructure constraints and other social and economic objectives for the district.
- 1.2.2 The Sustainability Appraisal for the District Plan has assessed a number of options for accommodating the level of growth required. The most sustainable strategy for the district is to focus development towards the three towns (Burgess Hill, Haywards Heath and East Grinstead) whilst allowing the villages to take growth to meet local housing and infrastructure needs. Strategic sites in Burgess Hill, have been deemed the most sustainable. Around 4,000 dwellings are planned for Burgess Hill, together with 20-30ha of employment land and supporting facilities and infrastructure. The District Council's preference is that the location and nature of development elsewhere in the district should be identified through the Neighbourhood Planning process. The District Plan will:
 - Put in place the overall planning framework for Mid Sussex with a coherent set of policies to protect and enhance the distinctive character of the District and its towns and villages for the next 17 years;
 - Encourage local communities to develop 'bottom up' neighbourhood plans; and

- Set out the Council's infrastructure needs and requirements in the Plan and ensure the necessary work on the Community Infrastructure Levy is completed in line with the Council's timetable.
- 1.2.3 Many of the District's Town and Parish Councils have started work on preparing their Neighbourhood Plans and some are well advanced in the statutory processes to get their plans in place. If it appears that the required level of development will not be delivered through Neighbourhood Plans, then the Council will need to consider producing its own allocations document.

1.3 Habitats Regulations Assessment

- 1.3.1 Habitats Regulations Assessment is a requirement of the Conservation of Habitats and Species Regulations 2010 (as amended; 'the Habitats Regulations'), the UK's transposition of *European Council Directive 92/43/EEC on the conservation of natural habitats and of wild fauna and flora* ('the Habitats Directive'). HRA must be applied to any plan or project in England and Wales with the potential to adversely affect the ecological integrity of any sites designated for their nature conservation importance as part of a system known collectively as the Natura 2000 network of European sites.
- 1.3.2 European sites provide ecological infrastructure for the protection of rare, endangered or vulnerable natural habitats and species of exceptional importance within the European Union. These sites consist of Special Areas of Conservation (SACs, designated under the Habitats Directive) and Special Protection Areas (SPAs, designated under *European Council Directive 2009/147/EC on the conservation of wild birds* ('the Birds Directive')). Meanwhile, the National Planning Policy Framework (DCLG, 2012) and Circular 06/05 (ODPM, 2005) require that Ramsar sites (UNESCO, 1971) are treated as if they are fully designated European sites for the purposes of considering development proposals that may affect them.

1.4 Structure of this Document

- 1.4.1 The outputs of the report include information in relation to:
 - Chapter 2 discusses the methodology for assessment, and updates the findings of the screening stage;
 - Chapter 3 defines relevant European sites, their qualifying features and conservation objectives;
 - Chapter 4 discusses the characteristics of European sites including available information about the population and ecology of qualifying species, and descriptions of qualifying habitats;
 - Chapter 5 presents available evidence regarding the mechanisms of atmospheric pollution impacts, and undertakes an assessment of these impacts in view of the sites' conservation objectives;

- Chapter 6 presents available evidence regarding the mechanisms of disturbance impacts, and undertakes an assessment of these impacts in view of the sites' conservation objectives;
- Chapter 7 presents a summary and concludes the document.

1.5 Relationship to Other Documents

- 1.5.1 This HRA Report draws on information previously published in other documents which form part of the overall HRA procedure for the Mid Sussex District Plan, including:
 - Habitats Regulations Assessment for the Mid Sussex District Plan: May 2013 (Urban Edge Environmental Consulting, 2013);
 - Ashdown Forest Visitor Survey Data Analysis (Natural England Commissioned Reports, Number 048: Clarke RT, Sharp J & Liley D 2010); and
 - Visitor Access Patterns on the Ashdown Forest: Recreational use and nature conservation (UE Associates and University of Brighton, 2009).



This page is intentionally blank.



2 Methodology

2.1 Guidance and Best Practice

- 2.1.1 Draft guidance on HRA has been defined by DCLG (2006) with more detailed draft guidance from Natural England (Tyldesley, 2009) and a range of other bodies¹. The guidance recognises that there is no statutory method for undertaking Habitats Regulations Assessment and that the adopted method must be appropriate to its purpose under the Habitats Directive and Regulations. DCLG guidance identifies three main stages to the HRA process:
 - Screening: Analysing draft options for likely significant effects on internationally designated sites;
 - Appropriate Assessment: Ascertaining the effects on site integrity; and
 - Alternative Solutions: Devising alternatives to the plan options, avoidance or mitigation measures.
- 2.1.2 An HRA must determine whether or not a plan or project will adversely affect the integrity of the European site(s) concerned, in view of the site's conservation objectives. Where adverse effects are anticipated changes must be made to the plan or project. The process is characterised by the precautionary principle. The European Commission (2000a) describes the principle as follows:

"If a preliminary scientific evaluation shows that there are reasonable grounds for concern that a particular activity might lead to damaging effects on the environment, or on human, animal or plant health, which would be inconsistent with the protection normally afforded to these within the European Community, the Precautionary Principle is triggered.

"Decision-makers then have to determine what action to take. They should take account of the potential consequences of taking no action, the uncertainties inherent in the scientific evaluation, and they should consult interested parties on the possible ways of managing the risk. Measures should be proportionate to the level of risk, and to the desired level of protection. They should be provisional in nature pending the availability of more reliable scientific data.

"Action is then undertaken to obtain further information enabling a more objective assessment of the risk. The measures taken to manage the risk should be maintained so long as the scientific information remains inconclusive and the risk unacceptable."

2.1.3 The hierarchy of intervention is important: where significant effects are likely or uncertain, decision-makers must firstly seek to avoid the effect through for example, a change of policy. If this is not possible, mitigation measures should be explored to remove or reduce significant

¹ For example European Commission (2001) and RSPB (Dodd et al, 2007)



effects. If neither avoidance, nor subsequent mitigation is possible, alternatives to the plan or project should be considered. Such alternatives should explore ways of achieving the objectives that avoid significant effects entirely. If there are no alternatives suitable for removing an adverse effect, decision-makers must demonstrate that there are Imperative Reasons of Overriding Public Interest to continue with the proposal. This is widely perceived as an undesirable position and should be avoided if at all possible.

2.2 Methodology

2.2.1 The guidance from Natural England (Tyldesley, 2009) was written for use in assessing strategic plans and forms the methodology adopted for this HRA; see Table 2.1. The overall objective of an Appropriate Assessment will be to ascertain whether any part of the plan will lead to an adverse effect on the ecological integrity of nearby European sites and, if so, make recommendations on how such effects can be avoided or mitigated.

2.3 Screening

2.3.1 An HRA screening assessment was prepared for the previous version of the District Plan, and this has been updated as the first step in preparing the current HRA. The screening assessment focuses the likely significant effects of each policy proposal on the European sites. Such effects can be sorted into one of 17 categories which are derived from the draft HRA guidance document produced for Natural England (Tyldesley, 2009). They help to determine which, if any, elements of the plan would be likely to have a significant effect on any interest feature of any European site, alone or in combination with other projects and plans, directly or indirectly. The 17 categories fall into four broader sections which are described as:

Category A	Elements of the plan / options that would have no negative effect on a European site at all
Category B	Elements of the plan / options that could have an effect, but the likelihood is there would be no significant negative effect on a European site either alone or in combination with other elements of the same plan, or other plans or projects
Category C	Elements of the plan / options that could or would be likely to have a significant effect alone and will require the plan to be subject to an appropriate assessment before it may be adopted
Category D	Elements of the plan / options that would be likely to have a significant effect in combination with other elements of the same plan, or other plans or projects and will require the plan to be subject to an appropriate assessment before the plan may be adopted

2.3.2 The categories, and traffic light colour-coded sub-categories, provide the means of recording the results of the assessment in such a way that important issues are identified whilst policies and proposed allocations that have no effect are screened out. Categories A, C and D are subdivided so that the specific reason why the assessor has allocated the policy or proposal to that category is more transparent, and more directly related to the ways in which the plan may affect a European site.



DCLG Stage	Natural England (Tyldesley) Ste	ps			
AA1: Likely significant effects	1. Gather the evidence base about international sites.				
	2. Consult Natural England and other stakeholders on the method for HRA and sites to be included.				
	3. Screen elements of the plans fo	r likelihood of significant effects.			
	4. Eliminate likely significant effect	ts by amending the plan / option.			
	5. Consult Natural England and other stakeholders on the findings of the screening stage, and scope of the Appropriate Assessment if required.				
AA2: Appropriate Assessment and ascertaining the effect on integrity		 8. Assess additions and changes to the plan and prepare draft HRA record. 9. Complete the draft 			
AA3: Mitigation measures and alternative solutions	7. Amend the plan / option or take other action to avoid any adverse effect on integrity of European site(s).	Appropriate Assessment and			
Reporting and	d 10. Submit draft HRA and supporting documents to Natural England.				
recording	11. Consult Natural England, other stakeholders and the public (if suitable).				
	12. Publish final HRA record and submit with Natural England letter to Inspector for Examination.				
	13. Respond to any representations relating to the HRA and to Inspector's questions.				
14. Check changes to the plan, complete HRA record and esta monitoring required.					

Table 2.1: Stages in HRA drawing on guidance from DCLG and Natural England

Findings of the screening stage

- 2.3.3 The screening assessment, which is presented at Appendix I, found that the following policy proposals were likely to significantly affect at least one European site:
 - DP2 Sustainable Economic Development;
 - DP3 Town Centre Development;
 - DP5 Housing;
 - DP7 Strategic Allocation to the east of Burgess Hill at Kings Way;
 - > DP8 Strategic Allocation to the north and north-west of Burgess Hill;
 - DP12 New Homes in the Countryside;



- DP29 Rural Exception Sites; and
- > DP30 Gypsies, Travellers and Travelling Showpeople.

2.4 Appropriate Assessment

2.4.1 The purpose of the Appropriate Assessment (HRA Stage AA2) is to further analyse likely significant effects identified during the screening stage, as well as those effects which were uncertain or not well understood and taken forward for assessment in accordance with the precautionary principle. The assessment should seek to establish whether or not the plan's effects, either alone or in combination with other plans or projects, will lead to adverse effects on site integrity, in view of the site's conservation objectives (see Chapter 3). Site integrity can be described as follows (ODPM, 2005):

"The integrity of a site is the coherence of its ecological structure and function, across its whole area, that enables it to sustain the habitat, complex of habitats and/or the levels of populations of the species for which it was classified."

2.5 In Combination Effects

2.5.1 Other plans and projects being prepared or implemented in the area may have the potential to cause negative effects on the integrity of European sites. These effects may be exacerbated when experienced in combination with the effects of the District Plan, possibly leading an insignificant effect to become significant. It is therefore important to consider which other plans and projects could generate similar effects as development within Mid Sussex, at the same European sites, and which may act in-combination. In combination effects are considered in Chapters 5 and 6.

3 European Sites

3.1 Scope of the Assessment

- 3.1.1 European sites considered within the scope of this assessment include all those falling partially within or close to Mid Sussex. Additionally, there may be activities occurring as a result of development within the district, which could take place outside of the confines of the district, possibly affecting European sites further afield.
- 3.1.2 The District Plan's forerunner, the draft Core Strategy, underwent an HRA screening and scoping exercise in late 2007 and early 2008 using a previous version of the Natural England guidance. This found that five designated areas were potentially at risk of effects from the Plan:
 - Ashdown Forest SAC;
 - Ashdown Forest SPA;
 - Castle Hill SAC;
 - Lewes Downs SAC; and
 - Mole Gap to Reigate Escarpment SAC.
- 3.1.3 Castle Hill, Lewes Downs and Mole Gap to Reigate Escarpment were screened out of the assessment, largely due to their distance from the district and the low likelihood of residents travelling along roads close to the sites. Natural England (2008) concurred with these findings in its screening opinion on the Core Strategy. Acknowledging that the plan is not necessary to the management of any European site, the screening exercise found likely significant effects on Ashdown Forest SAC/SPA as a result of disturbance and atmospheric pollution.
- 3.1.4 In accordance with Natural England's screening opinion, the assessment focuses on Ashdown Forest SAC/SPA; see Figure 3.1. These designations are described in the following sections.


3.2 Qualifying Features

3.2.1 The qualifying features of each site are listed in Table 3.1.

3.3 Conservation Objectives

3.3.1 The Habitats Directive requires that Member States maintain or where appropriate restore habitats and species populations of European importance to favourable conservation status. European site conservation objectives are referred to in the Habitats Regulations and Article 6(3) of the Habitats Directive. They are for use when there is a need to undertake an Appropriate Assessment under the relevant parts of the respective legislation.





Site	Qualifying Feature	Listing			
Ashdown	Primary reasons fo	r site selection			
Forest SAC (2,729 ha)	European dry heaths, for which this is considered to be one of the best areas in the United Kingdom.				
	Northern Atlantic wet heaths with <i>Erica</i> EC Habitats Directive 1992: Annex <i>tetralix</i> , for which this is considered to be Habitat one of the best areas in the UK.				
	Present but not a primary reason for site selection				
	Great crested newt <i>Triturus cristatus</i> , for which the area is considered to support a significant presence.				
Ashdown	Article 4.1 Qu	alification			
Forest SPA (3,207 ha)	Dartford warbler, 20 pairs representing 1.3% of the breeding population in Great Britain (Count, as at 1994).	EC Birds Directive 1979: Annex I			
	Nightjar, 35 pairs representing 1.0% of the breeding population in Great Britain (Count, as at 1991 and 1992).	EC Birds Directive 1979: Annex I			

Table 3.1: Qualifying features of the Ashdown Forest SAC and SPA

3.3.2 The conservation objectives are set for each feature (habitat or species) of an SAC/SPA. Where the objectives are met, the site can be said to demonstrate a high degree of integrity and the site itself makes a full contribution to achieving the aims of the Habitats and Birds Directives. Following the Habitats and Wild Birds Directives Implementation Review (HDIR; Defra 2012) Natural England committed to issuing a revised set of conservation objectives for the Natura 2000 network of SAC and SPA sites. The revised conservation objectives for Ashdown Forest SAC and SPA are listed in Table 3.2. Some of the typical species of each Annex 1 habitat are listed in Table 3.3. These are derived from a combination of sources, including the Joint Nature Conservation Committee (JNCC) Annex 1 habitat accounts and the Interpretation Manual of European Union Habitats (EC, 2007).

Table 3.2: Conservation objectives for Ashdown Forest SAC and SPA

Conservation Objectives for SAC

With regard to the natural habitats and/or species for which the site has been designated, the overriding conservation objective for each of the qualifying features is defined by Natural England as:

"Avoid the deterioration of the qualifying natural habitats and the habitats of qualifying species, and the significant disturbance of those qualifying species, ensuring the integrity of the site is maintained and the site makes a full contribution to achieving Favourable Conservation Status of each of the qualifying features."

Conservation objectives for the SAC are therefore, subject to natural change, to maintain or restore:

• Objective 1: The extent and distribution of qualifying natural habitats and habitats of qualifying species;

Conservation Objectives for SAC

- **Objective 2:** The structure and function (including typical species) of qualifying natural habitats and habitats of qualifying species;
- Objective 3: The supporting processes on which qualifying natural habitats and habitats of qualifying species rely;
- **Objective 4:** The populations of qualifying species;
- **Objective 5:** The distribution of qualifying species within the site.

Conservation Objectives for SPA

With regard to the individual species and/or assemblage of species for which the site has been classified, the over-riding conservation objective for each of the qualifying features is defined by Natural England as:

"Avoid the deterioration of the habitats of the qualifying features, and the significant disturbance of the qualifying species, ensuring the integrity of the site is maintained and the site makes a full contribution to achieving the aims of the Birds Directive."

Conservation objectives for the SPA are therefore, subject to natural change, to maintain or restore:

- **Objective 1:** The extent and distribution of the habitats of the qualifying features;
- **Objective 2:** The structure and function of the habitats of the qualifying features;
- > Objective 3: The supporting processes on which the habitats of the qualifying features rely;
- Objective 4: The populations of qualifying features;
- **Objective 5:** The distribution of qualifying features within the site.

Table 3.3: Typi	al species of Annex	1 habitat types prese	nt within SAC
-----------------	---------------------	-----------------------	---------------

Annex 1 Habitat Type	Typical Species
European dry heaths	Bell heather <i>Erica cinerea</i> , Dwarf gorse <i>Ulex</i> minor, Reptiles (adder, common lizard, sand lizard, smooth snake), Ants, bees and wasps (Hymenoptera), Beetles (Coleoptera), Dragonflies (Odonata)
North Atlantic wet heaths with <i>Erica tetralix</i>	Cross-leaved heath Erica tetralix, Sphagnum compactum, Deer grass Trichophorum cespitosum, Silver studded blue butterfly Plebejus argus

3.4 Condition Status

3.4.1 The conservation status of European sites is not routinely reported by Natural England, but it carries out condition monitoring of Sites of Special Scientific Interest (SSSI) at regular intervals. Although not exactly matching the boundaries of European sites, and being notified for different purposes, the condition status of a SSSI helps to give an impression of the overall ecological status of the SAC/SPA/Ramsar with which it coincides. The latest condition assessments of SSSIs forming part of the European sites within the scope of this assessment are summarised in Table 3.4.

% Eavourable		% Unfavourable no change		% Destroyed / part destroyed
15.69%	83.90%	0.00%	0.41%	0.00%



4 European Site Characterisation

4.1 Qualifying Species of the Special Area of Conservation

4.1.1 Although not a primary reason for site selection, great crested newt is present as a qualifying feature within the Ashdown Forest SAC.

Great crested newt

- 4.1.2 The great crested newt *Triturus cristatus* is the largest native British newt, reaching up to around 17cm in length. Adult males have jagged crests running along the body and tail. Newts require aquatic habitats for breeding. Eggs are laid singly on pond vegetation in spring, and larvae develop over summer to emerge in August October, normally taking 2–4 years to reach maturity. Great crested newt and their habitats are protected because the species has undergone significant declines over recent decades, primarily due to the loss of suitable breeding ponds as a result of agricultural intensification. Juveniles spend most time on land, and all terrestrial phases may range a considerable distance from breeding sites. The loss of terrestrial and connective habitats due to development has also been detrimental to the species' population.
- 4.1.3 The great crested newt is widespread throughout much of England and Wales, but occurs only sparsely in south-west England, mid Wales and Scotland. It is absent from Northern Ireland. The total UK population is relatively large and is distributed over sites that vary greatly in their ecological character. Furthermore, the UK has an internationally important population of great crested newts which have suffered serious decline throughout Europe. One estimate has put the national population at around 400,000 animals in 18,000 breeding sites. Many of the largest populations are centred on disused mineral-extraction sites, but lowland farmland forms the majority of great crested newt habitat in the UK.

4.2 Qualifying Habitats of the Special Area of Conservation

4.2.1 The following sections are adapted from the JNCC site accounts of the Ashdown Forest SAC.

European dry heaths

- 4.2.2 Ashdown Forest contains one of the largest single continuous blocks of lowland heath in southeast England, with both European dry heaths and, in a larger proportion, North Atlantic wet heath.
- 4.2.3 The dry heath in Ashdown Forest is an extensive example of the south-eastern Calluna vulgaris
 Ulex minor community. This vegetation type is dominated by heather Calluna vulgaris, bell heather Erica cinerea and dwarf gorse Ulex minor, with transitions to other habitats. It supports



important lichen assemblages, including species such as *Pycnothelia papillaria*. This site supports the most inland remaining population of hairy greenweed *Genista pilosa* in Britain.

Northern Atlantic wet heaths with Erica tetralix

4.2.4 The Erica tetralix – Sphagnum compactum wet heath element provides suitable conditions for several species of bog-mosses Sphagnum spp., bog asphodel Narthecium ossifragum, deergrass Trichophorum cespitosum, common cotton-grass Eriophorum angustifolium, marsh gentian Gentiana pneumonanthe and marsh clubmoss Lycopodiella inundata. The site supports important assemblages of beetles, dragonflies, damselflies and butterflies, including the nationally rare silver-studded blue Plebejus argus, and birds of European importance, such as European nightjar Caprimulgus europaeus, Dartford warbler Sylvia undata and Eurasian hobby Falco subbuteo.

4.3 SPA Bird Populations

- 4.3.1 Ashdown Forest is located in the High Weald of East Sussex in south-east England, where valley mires, heath and damp woodland have developed on soils derived from Hastings Sands (Lower Cretaceous). Once a royal hunting forest, reduced grazing has resulted in the accelerated development of woodland and encroachment of bracken *Pterdium aquilinium* over former heath. Nevertheless, some fine examples of heathland habitats remain, with humid or wet heath predominating (around 45% cover), dominated by heather, bell heather and cross-leaved heath *E. tetralix* in the dampest conditions. Where drier heaths occur (around 15% cover) they are dominated by heather in association with gorse *Ulex europaeus* and dwarf gorse. Streamsides and mires add further variety (around 5% cover), with *Sphagnum* mosses, cottongrass *Eriophorum* sp., bog asphodel and round-leaved sundew *Drosera rotundifolia* all characteristic plants. The woodlands (around 35% cover) are also varied, with birch *Betula* sp. typically establishing first over heath, followed by oak *Quercus robur*, willow *Salix* sp. and pine *Pinus* sp. in places, eventually forming dense and shaded areas with sparse ground flora.
- 4.3.2 Together with the nearby Wealden Heaths SPA and Thames Basin Heath SPA, Ashdown Forest forms part of a complex of heathlands in southern England that support breeding bird populations of European importance in particular nightjar *Caprimulgus europaeus* and Dartford warbler *Sylvia undata*. Breeding birds of scrub and woodland (such as woodlark *Lullula arborea* [also an Annex 1 species occurring in qualifying numbers] and Eurasian hobby *Falco subbuteo*) are also associated with the varied mosaic of their respective habitats, distributed over the higher slopes and valleys of the High Weald.

Dartford warbler

4.3.3 Although native to the UK, the Dartford warbler is really a species of the Mediterranean. The European population is declining at a moderately rapid rate, qualifying the species as Near Threatened (IUCN, 2012). Declines in the core population in Spain are largely responsible for overall declines. The drivers of this decline are not entirely clear but include habitat degradation and modification.

4.3.4 In the UK, a national survey in 1994 recorded 1,889 breeding territories (Gibbons & Wotton, 1996), a recovery from the population crash of the 1960s following cold winters. A repeat survey in 2006 showed that the national population had increased with 3,214 territories recorded (Wotton *et al.*, 2009). Dartford warbler is Amber listed on Birds of Conservation Concern 3 (BoCC3; Eaton *et al*, 2009). There were 38 territories recorded within the Ashdown Forest SPA during the 2006 survey (Wotton *et al.*, 2009).

Nightjar

- 4.3.5 As reported by Baillie *et al*, 2012, after a substantial range contraction of more than 50% of 10km squares up to 1981, the 1992 national survey revealed a 50% increase in population size since 1981, probably due to the increased availability of young forest habitat as plantations were felled and replanted (Morris *et al.* 1994).
- 4.3.6 A National Nightjar Survey in 2004 revealed that a further 36% increase had taken place in the UK population in 12 years, with a 2.6% increase in the number of 10km squares occupied (Conway et al. 2007). However, a steep linear decrease in the number of fledglings per breeding attempt has been observed through the BTO's nest recording scheme (Baillie et al., 2012). A recent study suggests that nest failure is most likely in areas heavily frequented by walkers and dogs (Langston et al. 2007). Nightjar is Red listed on BoCC3. A total of 83 churring males were recorded within the SPA during the 2004 national survey (Conway et al. 2007).

Woodlark

- 4.3.7 As reported by Baillie *et al*, 2012, woodlark suffered a 62% range contraction between 1968-72 and 1988-91; the species had ceased to breed in Wales and in several southern English counties over this period (Gibbons *et al.* 1993). Sitters *et al.* (1996) found that the UK population increased from c.250 pairs in 1986 to c.600 pairs in 1993, probably helped by mild winters and increased habitat availability due to storm damage in plantations, forest restocking, and heathland management. A repeat national survey in 1997 showed that the population had increased further, accompanied by expansion of the range into new areas (Wotton & Gillings 2000). A further repeat in 2006 recorded an increase since 1997 of 88% accompanied by major range expansion, with a pair breeding in Wales for the first time since 1981 (Conway *et al.* 2009).
- 4.3.8 Farmland set-aside, especially close to forest, was valuable additional habitat for the expanding population, although clutch sizes may be lower there than in more traditional habitats (Wright *et al.* 2007). Climate change may benefit woodlark, because it is able to make more nesting attempts in warmer years (Wright *et al.* 2009). The small nest record scheme sample suggests that nest failure rates have become less frequent at the egg stage, but there has been no trend in the number of fledglings per breeding attempt. Human disturbance at heathland sites apparently reduces population density, but the effects are partly offset by higher breeding productivity at lower densities (Mallord *et al.* 2007). The species' partial recovery in numbers and range resulted in a move from the Red to the Amber list at the 2009 BoCC review (Eaton *et al.* 2009). A total of 42 territories were recorded within the SPA during the 2006 national survey (Conway *et al.* 2009).



This page is intentionally blank.

5 Atmospheric Pollution Impact Pathways

5.1 Introduction

- 5.1.1 The screening exercise found that the following District Plan policies were likely to significantly affect Ashdown Forest SAC as a result of atmospheric pollution impacts:
 - DP2 Sustainable Economic Development;
 - DP3 Town Centre Development;
 - DP5 Housing;
 - > DP7 Strategic Allocation to the east of Burgess Hill at Kings Way; and
 - > DP8 Strategic Allocation to the north and north-west of Burgess Hill.

5.2 Understanding the Vulnerability of SAC Habitats to Atmospheric Pollution

- 5.2.1 Atmospheric pollution is a widespread issue, with background air quality heavily influenced by large point-source emitters including transboundary sources. Local pollutant sources can affect designated sites, particularly in relation to protected habitats within SACs, and especially from road traffic emissions. The District Plan cannot feasibly influence causes of background pollution such as large point sources but, through the scale and distribution of development proposed and sustainable transport measures, will affect the way in which locally emitted pollutants reach the site.
- 5.2.2 Qualifying habitats most sensitive to air pollution within Ashdown Forest are European dry heaths and North Atlantic wet heaths. The main pollutant effects of interest are acid deposition and eutrophication by nitrogen deposition. The following brief descriptions draw on information presented through the Air Pollution Information System² (APIS).
- 5.2.3 Acid deposition: caused by oxides of nitrogen (NO_x) (or sulphur dioxide) reacting with rain/cloudwater to form nitric (or sulphuric) acid, and is caused primarily by energy generation, as well as road traffic and industrial combustion. Both wet and dry acid deposition have been implicated in the damage and destruction of vegetation (heather, mosses, liverworts and lichens are particularly susceptible to cell membrane damage due to excessive pollutant levels) and in the degradation of soils and watercourses (including acidification and reduced microbial activity).
- 5.2.4 Eutrophication by nitrogen deposition: consists of the input of nitrogen from NO_X (and sometimes ammonia) emissions by deposition, and is caused primarily by road traffic, as well as energy generation, industrial combustion and agricultural practices. Nitrogen deposition can cause direct damage to heather, mosses, liverworts and lichens, as well as other plant species,

²Online at: <u>http://www.apis.ac.uk/index.html</u> [Accessed 25/7/11]



because of their sensitivity to additional atmospheric nitrogen inputs, whilst deposition can also lead to long term compositional changes in vegetation and reduced diversity. For example a marked decline in heather and an increased dominance of grasses have been observed throughout the Netherlands and also in the East Anglian Brecklands (see for example Bobbink et al (1993) and Pitcairn et al (1991)).

- 5.2.5 Furthermore, while plants are able to detoxify and assimilate low exposure to atmospheric concentrations of NO_X, high levels of uptake can lead to detrimental impacts including:
 - > Inhibition of pigment biosynthesis, leading to reduced rates of photosynthesis;
 - Water soaking as NO₂ molecules attach to lipids in membranes, causing plasmolysis (removal of water) and eventually necrosis;
 - > Inhibition of lipid biosynthesis, leading to reduced rates of regeneration and growth;
 - > Injury to mitochondria and plastids, essential to internal processing of energy & proteins;
 - > Decrease in stomatal conductance of air and water vapour; and
 - Inhibition of carbon fixation (at least under low light levels).
- 5.2.6 Over half of all emissions of nitrogen and nitrogen oxides in the UK are the result of vehicle exhausts, with an estimated 92% of those associated with residential development being contributed by road traffic (Dore *et al*, 2005). Nitrogen emissions from traffic generated by residential and commercial developments will therefore be the focus of this part of the assessment. The scope can be further refined by concentrating on traffic growth on roads within 200m of the SAC, as beyond 200m effects of emissions from this source diminish to the equivalent of background levels (Laxen & Wilson (2002)).

5.3 Evidence for Pollution Impacts within Ashdown Forest

- 5.3.1 The critical load for nitrogen deposition and acid (nitrogen (N) or sulphur (S)) deposition is already exceeded in parts of Ashdown Forest; Table 5.1 presents information on background critical load exceedances for these key pollutants on qualifying habitat types at a selection of grid references across the Forest, shown on Figure 5.1. Critical loads concern the quantity of pollutants deposited from the air to the ground. Nilsson and Grennfelt (1988) define critical loads as "a quantitative estimate of exposure to one or more pollutants below which significant harmful effects on specified sensitive elements of the environment do not occur according to present knowledge".
- 5.3.2 Although a nitrogen deposition critical load range is expressed for each habitat in the below table, APIS gives further guidance on critical loads to be used when assessing nitrogen deposition in impact assessments. This confirms that the lower end of the range should be used at both the screening stage and detailed assessment stage when assessing impacts on northern wet heath and dry heath habitats (and broadleaved woodland).



Receptor	N Deposition (kg/ha/yr)		Acid dep (keq N/ha/yr)		Acid dep (keq S/ha/yr)	
	Crit. load	Actual load	Crit. load	Actual load	Crit. load	Actual load
1. Woodland	10-20	28.84	0.14-3.23	2.06	3.08	0.31
2. Dry heath	10-20	15.96	0.64-0.95	1.31	0.31	0.27
3. Wet heath	10-20	16.66	0.64-0.97	1.31	0.33	0.27

Table 5.1: Critical load for Ashdown Forest SAC compared to actual maximum load

1. Nearest SAC location to East Grinstead and close to A22; woodland habitat (not designated). NGR542021,133634. APIS Habitat: Broadleaved, mixed and yew woodland (N Dep.) / Broadleafed/coniferous unmanaged woodland (Acid Dep.). Data are 2010-2012.

2. Dry heath habitat location close to A22 at Ashdown Llama Farm. NGR542234,131385. APIS Habitat: Dry heaths (N. Dep.) / Dwarf shrub heath (Acid Dep.). Data are for the period 2009-2011.


3. Wet heath habitat location close to A22 at Millbrook. NGR544045,128936. APIS Habitat: Northern (*E. tetralix* dominated) wet heath (N Dep.) / Dwarf shrub heath (Acid Dep.). Data are for the period 2009-2011.

5.3.3 The data in Table 5.1 is historical (2011) and provides an indicative assessment as to which areas are approaching the limits of environmental capacity. In all cases, the critical load for nutrient nitrogen and acid nitrogen deposition is exceeded, indicating that significant additional sources of these pollutants generated as a result of proposals in the District Plan should be avoided or mitigated to prevent additional adverse effects on ecological integrity. The following section establishes whether the District Plan is likely to contribute significant additional pollution deposition.

5.4 Contribution of the District Plan to Future Pollution Deposition within Ashdown Forest

- 5.4.1 The Design Manual for Roads and Bridges (DMRB; Highways Agency, 2007) provides guidance on assessment of the impact that road projects may have on local air quality. Specific provision is made in relation to sites designated under the Habitats Directive. In this instance the assessment is in relation to existing, as opposed to new roads, however the guidance clarifies that 'where appropriate, the advice may be applied to existing roads'. In accordance with this guidance, and with agreement from Natural England (minutes of meeting between Natural England, Mid Sussex District Council and Wealden District Council, 16 September 2010), the HRA examines whether there is a likely significant effect using the DMRB guidance.
- 5.4.2 DMRB provides a scoping assessment for local air quality and initially requires the identification of roads which are likely to be affected by the proposals. The criteria for defining an affected road are:
 - Road alignment will change by 5 metres or more; or
 - > Daily traffic flows will change by 1,000 annual average daily traffic (AADT) or more; or
 - Heavy Duty Vehicle (HDV) flows will change by 200 AADT or more; or
 - > Daily average speed will change by 10km/hr or more; or
 - Peak hour speed will change by 20km/hr or more.
- 5.4.3 The scoping assessment then requires that nature conservation sites (e.g. SACs) within 200m of the road and their characteristics be identified.





5.4.4 The guidance clarifies that if none of the roads in the network meet the traffic/alignment criteria (that is, they are not affected roads) or there are no relevant designated sites near the affected roads, then the air pollution impact of the scheme is considered not to be significant and no further work is needed. Major roads passing through the Forest along which residents from Mid Sussex could be likely to travel are: A22, A26 and A275, while the B2188, B2026, B2110 and Coleman's Hatch Road may also be of concern; see Figure 5.2.


Mid Sussex Transport Study Report (December 2012)

- 5.4.5 Mid Sussex District Council commissioned Amey Transport Consultants to carry out the Mid Sussex Transport Study Stage 1 (December 2012) in order to assess and understand the transport impacts of its policies and proposals in the Mid Sussex District Plan. Part of the brief required an assessment of whether or not future land use and travel patterns in Mid Sussex would impact on the environmentally sensitive habitats of the Ashdown Forest SAC.
- 5.4.6 The study utilises a derivative of the West Sussex County Council strategic multi modal transport model (WSCCM) to assess the transport impacts of the District Plan. Using the Design Manual for Roads and Bridges guidance, representative local flow factors were applied to convert forecast AM peak model outputs at 2021 and 2031 to AADT equivalent.

Study area

- 5.4.7 The transport model was specifically refined to include new network detail covering roads that access or cross Ashdown Forest, namely:
 - A275 (Lewes-East Grinstead);
 - A22 (Uckfield-East Grinstead);
 - A26 (Uckfield-Crowborough);
 - B21110 (East Grinstead-Royal Tunbridge Wells);
 - B2110 (East Grinstead- Royal Tunbridge Wells);
 - B2188 (Maresfield Groombridge);
 - B2026 (B2188-B2110); and
 - Coleman's Hatch Road (East –West through Ashdown Forest).
- 5.4.8 Owing to the strategic nature of the Mid Sussex Transport Study and the location of Ashdown Forest on the north east periphery of the modelled network, meaningful flow assignments for B2188, B2026 and Coleman's Hatch Road could not be derived. However an assessment has been made of future traffic impacts on A275, A22, A26 and B2110 which are the main routes that are used in the area and experience most traffic flows.





Treatment of development scenarios

- 5.4.9 The District Plan identifies the overall amount of new development needed over the period 2014-2031 and identifies strategic allocations at Burgess Hill. The Council has produced an estimate of objectively assessed housing needs in Mid Sussex to inform the decisions made regarding the levels of growth that will be appropriate in the district. This number is currently 516 per annum (8,776 to 2031) although the Plan requirement number is still subject to confirmation. The introduction of Neighbourhood Planning has also influenced the strategy contained within the District Plan.
- 5.4.10 The District Plan policies for strategic development at Burgess Hill require approximately 4,000 new homes to the north/north-west and east of the town, together with up to 20 30 hectares of employment land as a high quality business park and sporting facilities. The residential development is identified as two strategic development allocations of land to the east of Burgess Hill at Kings Way and land to the north and northwest of Burgess Hill.
- 5.4.11 The Council's preference is that the location and nature of the remaining homes and employment land are identified through Neighbourhood Plans. However for the work of the Transport Study a number of Development Scenarios were tested which included proposed strategic sites at Burgess Hill and housing likely to come forward elsewhere which was distributed proportionally, according to defined 'Zones'. These are Zones based on urban and rural areas. Further detail is provided in Chapter 3 of the Mid Sussex Transport Study Report.

Results

- 5.4.12 The findings show that the maximum two-way AADT flow increase on A275 in 2021, will be 179 vehicles in the scenario without remedial transport interventions and 100 vehicles with transport interventions. Similarly the maximum two way AADT flow on A275 in 2031 will be 417 vehicles (without remedial transport interventions) and 358 (with transport interventions). The suggested transport interventions are set out in the Mid Sussex Transport Study Report Stage 1, which concludes that traffic impacts will be less on A22, A26 and B2100 than they are predicted to be on A275 (refer to Appendix C of the transport study for further details).
- 5.4.13 All projected traffic increases are therefore expected to fall well below the 1,000 AADT increase threshold, specified in the Design Manual for Roads and Bridges Guidance, as the level below which air pollution impacts from road traffic can be considered not significant in relation to European sites.

5.5 Other Plans and Projects acting In Combination

5.5.1 Other plans and projects which may act in combination with the District Plan to exacerbate pollution impacts to the SAC include developments proposals within Wealden district. Wealden District Council is implementing its adopted Core Strategy Local Plan, which includes a spatial distribution of development which was demonstrated not to result in significant traffic-related air pollution impacts on Ashdown Forest. In assessing traffic flow increases on roads close to Mid Sussex within the SAC, the HRA for Wealden's Core Strategy (UE Associates, August 2011, p.25) predicts AADT traffic growth of 583 vehicles on the A22 north of Forest Row.



It does not include predictions for the A275 which is the only road for which figures are available from the Mid Sussex Transport Study. However, the latter concluded that traffic growth on the A22 resulting from Mid Sussex developments would be less than predicted on the A275 (358 vehicles in 2031 with transport interventions), meaning that the combined traffic flow on the A22 is not predicted to exceed the 1,000 AADT significance threshold for European sites.

5.5.2 Wealden District Council's website³ includes the following statement:

"In addition to recreational impacts, the Council is also obliged to assess all development which would, either alone or in combination with other development, increase the level of nitrogen deposition on the Ashdown Forest SAC. When considering nitrogen deposition the focus will be on the emissions from motorised vehicular traffic... Where it is assessed that there will be a likely significant effect due to an increase in the level of nitrogen deposition, and in the absence of any identified mitigation measures, planning permission must be withheld."

5.5.3 Wealden District Council has also embarked on a long-term monitoring and research project to provide information on the impacts of nitrogen deposition on the Ashdown Forest SAC so that its effects on development in the longer term can be more fully understood. The requirement to implement the monitoring programme was one of the recommendations of the independent Planning Inspector who assessed the Wealden's Core Strategy as sound.

5.6 Avoidance and Mitigation Measures proposed by the District Plan

5.6.1 The District Plan contains measures to promote sustainable transport over the plan period, including measures relating to existing development, and additional actions to assess and manage air pollution. These are intended to improve the overall sustainability of the District as well as reduce the traffic emissions from proposed development, including along roads passing through or close to Ashdown Forest; see Box 1.

Box 1: Policy DP18 – Transport

Development will be required to support the objectives of the West Sussex Local Transport Plan, which are:

- A high quality transport network that promotes a competitive and prosperous economy;
- A resilient transport network that complements the built and natural environment whilst reducing carbon emissions over time;
- Access to services, employment and housing; and
- A transport network that feels, and is, safer and healthier to use.

To meet these objectives, development will only be permitted where:

- It is sustainably located to minimise the need for travel;
- It facilitates and promotes the increased use of alternative means of transport to the private

³ Wealden District Council website [accessed September 2014]: http://www.wealden.gov.uk/Wealden/Residents/Planning_and_Building_Control/Planning_Development_Management/Agents_and_Parish_Council_Information/Planning_Agents_Ashdown_Forest.aspx



car, such as the provision of, and access to, safe and convenient routes for walking, cycling and public transport;

- Does not cause a severe impact in terms of road safety and increased traffic congestion;
- Is designed to adoptable standards, or other standards as agreed by the Local Planning Authority, including road widths and size of car parking spaces and size of garages; and
- Provides adequate car parking for the proposed development in accordance with parking standards as agreed by the Local Planning Authority or in accordance with the relevant Neighbourhood Plan.

Where practical developments should be located and designed to incorporate facilities for charging plug-in and other ultra-low emission vehicles.

Depending on the size of and the likely transport impact of development, a Transport Statement or Transport Assessment will be submitted alongside planning applications.

The District Council will continue to work with public transport providers to ensure that the level of service provided meets the needs of commuters and other travellers. This includes the frequency of services and quality of infrastructure such as stations and trains.

Neighbourhood Plans can set local standards for car parking provision provided that it is justified by evidence.

5.7 Impact Assessment

5.7.1 This section assesses whether the District Plan can be expected to adversely affect the ecological integrity of the Ashdown Forest SAC as a result of atmospheric pollution impacts. It does this by considering the sites' conservation objectives (Table 3.2), with reference to the characteristics of the site (Chapter 4) and its qualifying features (Table 3.1), and in the context of available data on pollution impacts and the District Plan's policy response as outlined above (sections 5.1 to 5.6).

Objective 1: Extent and distribution of qualifying natural habitats and habitats of qualifying species

5.7.2 Ashdown Forest is designated as an SAC for its European dry heaths and North Atlantic wet heaths with *E. tetralix*. Great crested newts are present in qualifying numbers but are not a primary reason for site selection. Predicted traffic growth as a result of development proposed by the District Plan is below the significance threshold set for European sites. The extent and distribution of qualifying natural habitats and habitats of qualifying species are unlikely to be affected.

Objective 2: Structure and function (including typical species) of qualifying natural habitats and habitats of qualifying species

5.7.3 Predicted traffic growth is below the significance threshold set for European sites; the structure and function of qualifying natural habitats and habitats of qualifying species are unlikely to be affected.



Objective 3: Supporting processes on which qualifying natural habitats and habitats of qualifying species rely

5.7.4 Predicted traffic growth is below the significance threshold set for European sites; the supporting processes on which qualifying natural habitats and habitats of qualifying species rely are unlikely to be affected.

Objective 4: Populations of qualifying species

5.7.5 Predicted traffic growth is below the significance threshold set for European sites; the populations of qualifying species are unlikely to be affected.

Objective 5: Distribution of qualifying species within the site

5.7.6 Predicted traffic growth is below the significance threshold set for European sites; the distribution of qualifying species within the site is unlikely to be affected.

5.8 Conclusions regarding Adverse Effects on Integrity

5.8.1 It can be concluded that the District Plan will not result in adverse effects on the ecological integrity of the Ashdown Forest SAC.



6 Disturbance Impact Pathways

6.1 Introduction

- 6.1.1 The screening exercise found that the following District Plan policies were likely to significantly affect Ashdown Forest SPA as a result of disturbance impacts:
 - DP5 Housing;
 - DP12 New Homes in the Countryside;
 - DP29 Rural Exception Sites; and
 - > DP30 Gypsies, Travellers and Travelling Showpeople.

6.2 Understanding the Vulnerability of SPA Species to Disturbance

6.2.1 Murison *et al.* (2007) noted that animals often react to human disturbance as a form of predation risk. Such a response can include elevated heart rate, heightened defensive behaviour, including evasive measures with associated energy expenditure, and the avoidance of high risk areas (Murison *et al.* (2007), Liley & Sutherland (2007)). High levels of human activity in important nature conservation areas might then change the behaviour of animals to such a degree that conservation priorities become compromised. This may result from reduced breeding success, increased predation or exposure of nests, eggs or young to trampling and the elements (Liley & Sutherland, 2007).

Dartford warbler

- 6.2.2 In a study into the relationship between habitat type and disturbance effects on the breeding Dartford warbler, Murison *et al.* (2007) noted the following as important measures of disturbance. First, they noted that indirect disturbance was associated with factors such as the distance from the centre of the heathland patch (or nest) to the nearest road, path, building or car park. Second, the proximity of a nest territory to the nearest access point showed a strong, negative relationship with the timing of a first brood. Third, disturbance appeared to be associated with increased stress levels, with birds exhibiting an extended period of agitation while searching for cover, leading to increased energy expenditure.
- 6.2.3 They suggest that the mechanisms by which disturbance affects the Dartford warbler's breeding success are associated with its particular susceptibility to disturbance during nest-building, with birds often abandoning their work and materials. The effects of this are threefold. The timing of the first brood was delayed for long enough (up to six weeks) to prevent multiple broods in one season. Second, the fledgling success of a first brood delayed until June was limited by the decreased availability of invertebrate prey. Finally, that continued disturbance events reduced the foraging effectiveness of the birds, and their ability to feed their young, by keeping the adults away from the nest for longer than normal.



6.2.4 Analysing the results of their study, Murison *et al.* (2007) found that breeding pairs with territories in areas experiencing as many as 13 to 16 disturbance events each hour of every day, delayed breeding for sufficiently long enough to prevent multiple broods in one season. Importantly, they also found a significant correlation between the reproductive success of Dartford warbler and the proportion of different species of gorse present in the heathland patch. They discovered a strong positive relationship with European gorse *Ulex europaeus*, where heathland patches containing more of this type produced more successful broods. While the significance of disturbance events in delaying breeding among pairs nesting in heather-dominated territories was high, often leading to reduced breeding success, the correlation was weaker in territories dominated by Western gorse *U. gallii*. During their surveys, dogs were observed ranging as far as 45m into heather dominated areas, but never strayed from the path in areas with vegetation dominated by gorse.

Nightjar

- 6.2.5 Liley and Clarke (2003), following field studies into the population density of nightjar on 36 patches of heathland in Dorset, demonstrated that patches surrounded by higher levels of development supported smaller populations of nightjar. Effects associated with urbanisation identified as relevant in this respect included human disturbance, light pollution, predation from natural predators and domestic pets (as well as corvids, Fox *Vulpes vulpes*, and Hedgehog *Erinaceus europaeus*), and habitat change.
- 6.2.6 In a study investigating the relationship between walkers with dogs and the success of breeding nightjar, Langston *et al.* (2007) observed that the flushing of birds from the nest by a disturbance event during daylight hours led to predation by diurnal predators, particularly of eggs. Moreover, birds tend to flush more readily in response to dogs than to humans, and take longer to return to the nest. Langston *et al.* (2007) noted that disturbance effects on nightjar were accentuated when breeding conditions were less favourable due to incidental factors such as weather conditions. Birds flushing the nest as a result of disturbance events during harsh or wet weather tended to bear smaller, less successful broods. Overall, they found a significant relationship between nest failure and disturbance, with failure being more likely in nests with higher total footpath length within 50m, 100m and 500m of the nest clearing.

Woodlark

- 6.2.7 It has been observed that the removal of human disturbance effects could result in an increase of between 13% and 48% in the breeding population of woodlark over 16 heathland sites (Mallord *et al.* 2007a, Mallord *et al.* 2007b). At sites with recreational access woodlark was found to be less likely to colonise suitable habitat in areas with greater disturbance. The probability of colonisation was reduced to below 50% with disturbance levels at eight events per hour.
- 6.2.8 In summary, disturbance impacts to heathland breeding birds can be described as:
 - Increased nest predation by natural predators or dogs when adults are flushed from the nest or deterred from returning to it by the presence of people or dogs;
 - Chicks or eggs dying of exposure because adult birds are kept away from the nest;

- Accidental trampling of eggs by people, where nests are on the ground and may be close to paths;
- Reduced productivity resulting from delayed breeding, fewer young, decreased food availability and/or nest failure;
- Displacement from / avoidance of otherwise suitable areas of habitat; and
- Increasing stress levels in adult birds in response to perceived predation risk.

6.3 Evidence for Disturbance Impacts within Ashdown Forest

- 6.3.1 The study by Clarke *et al.* (2010), analysed data on visiting patterns collected through face-toface questionnaire surveys conducted at Ashdown Forest in 2008 (UE Associates & University of Brighton, 2009). After assessing visitor rates at each access point, they distributed visitor numbers across the path network within the Forest using a 25m by 25m cell grid to generate a series of maps showing visitor intensity levels. These were overlaid with recorded bird territories from the latest national surveys for Dartford warbler (Wotton *et al.*, 2009), nightjar (Conway *et al.* 2007) and woodlark (Conway *et al.* 2009) to investigate whether bird density correlates with visitor or path intensity.
- 6.3.2 They found that average bird density was lowest in areas with the lowest two class levels of either visitor or path intensity. For woodlark, the highest average density occurs in areas with the highest class levels of visitor intensity at all three assessed influence distances (50m, 100m and 150m). Average nightjar density was lowest for the quarter of cells with the lowest nearby visitor or path intensity for all three influence distances. Dartford warbler average density was highest amongst the quarter of cells with the highest level of visitor intensity. For each species, several of the differences in bird density with nearby visitor or path intensity were statistically significant (Clarke *et al.* 2010). The positive relationship between nesting birds and presence or potential presence of visitors would seem to suggest that visitor activity is not affecting territory selection among the birds on Ashdown Forest.
- 6.3.3 However, after accounting for habitat type, Clarke *et al.* (2010) found that path intensity is lowest in (or very near to) areas of wet heath, and second lowest in woodland. Furthermore, visitor intensity levels were found to be much lower in areas of wet heath and woodland than other habitats, and much less than half the general level of visitor pressure in area of dry heath. Looking at the habitat preferences of the birds within Ashdown Forest, wet heath and woodland habitats are less preferred by the bird species, whereas dry heath was found to be the preferred nesting habitat for Dartford warbler and nightjar, and second choice (of six habitat types) among woodlark. Other important nesting habitats were gorse (especially for woodlark and Dartford warbler) and unimproved grassland where levels of visitor and path intensity were also higher than in woodland and wet heath habitats.
- 6.3.4 Analysis which also takes account of habitat type thus demonstrates that more open and drier habitats are generally preferred by both birds and people within Ashdown Forest, perhaps explaining why the density of nesting territories is greatest in areas with the highest presence or potential presence of visitors. But Clarke *et al.* (2010) found no clear evidence that the current spatial distributions of woodlark, nightjar or Dartford warbler are affected by the patterns of



current levels of nearby visitor pressure or by path intensity within the SPA. Nevertheless, the density of all three species is low within Ashdown Forest, and lower than might be expected when compared to other heathland sites in southern England with a similar range of habitats (see Clarke *et al.*, 2010, Table 7 and Figure 14). By comparison, visitor pressure at the site (2.1767 visitors per ha over a 16 hour period) appears to be lower than on the Thames Basin Heaths (3.7 visitors per ha) and slightly higher than the Dorset Heaths (1.8 visitors per ha) where similar studies have been carried out.

6.3.5 The study concludes that the current (2008) level of visitor pressure is not affecting the distribution of nightjar, woodlark or Dartford warbler within Ashdown Forest SPA; based on the analysis undertaken, the birds do not appear to be avoiding areas of greater recreational pressure. However, recreational disturbance could still be having an impact on the Annex I bird species at the site:

"It may be that the density of birds is so low (due to other, currently unknown factors) that there is little competition for space and therefore no impacts of disturbance. While the results of the analysis presented in the report are potentially encouraging, in the absence of data on breeding success, and without understanding why bird densities are low, it currently cannot be concluded on the basis of scientific evidence that the ecological integrity of nightjar and Dartford warbler populations is not being adversely affected by a combination of existing pressure and/or habitat management." Clarke et al., 2010, p29

6.3.6 Clarke *et al.* (2010) advocate a precautionary approach when considering the implications of a plan or project proposing a significant level of increased housing, informed by wider research into the effects of disturbance on heathland birds, because it cannot be conclusively determined that current levels of recreational pressure are not affecting the breeding success of birds exposed to recreational pressure.

6.4 Determining a Zone of Influence for Visiting Patterns to Ashdown Forest

- 6.4.1 Using the 2008 visitor data, Clarke *et al.* (2010) estimated that a total of 5,198 people would currently visit Ashdown Forest over a 16 hour period in September, or 325 visits per hour. Given the visitable area of the SPA (2,388ha), this equates to a visitor density of 2.17 visitors per ha over 16hrs. In order for such information to be useful for spatial planning purposes, it is necessary to define how far people will travel from home to visit Ashdown Forest, and thus where new housing development is most likely to increase visitor pressure in the SPA such that impacts to nesting heathland birds could occur.
- 6.4.2 For example, development where residents are likely to visit Ashdown Forest on a daily basis to walk the dog (i.e. those closest to the Forest) should be considered differently from development further away where residents might visit Ashdown Forest more occasionally. Research (for example UE Associates and University of Brighton (2009), Underhill-Day & Liley (2007), Underhill-Day (2005)) and practical experience in protecting the Thames Basin Heaths SPA and Dorset Heathlands SPA has shown that avoiding or mitigating the disturbance impacts of residential development within 400m of the SPA is unlikely to be successful, and hence that residential development within this distance should not be permitted.



6.4.3 In both the Dorset and Thames Basin Heaths a 5km zone has been established around the SPA. Within this zone it is recognised that new development has the potential to result in significantly increased use of the heaths, and that mitigation measures are required to reduce this increase, funded through developer contributions. Figure 6.1 shows the cumulative percentage of visitors to Ashdown Forest arriving by car in relation to distance from the Forest, and compares this to similar patterns at the Thames Basin and Dorset Heaths. It shows that visitors to Ashdown Forest appear to travel further than visitors to the Dorset or Thames Basin Heaths. For instance, around 40% of interviewed visitors travelling by car to Ashdown Forest had come from within a 5km distance. On the Thames Basin Heaths, this figure was approximately 60% and at the Dorset Heaths it was around 70%. At Ashdown Forest, approximately 65% of visitors arriving by car travel from within 7km of the site.




Figure 6.1: Cumulative frequency of car visitors by distance, for the Dorset Heaths, Thames Basin Heaths and Ashdown Forest (Source: Clarke et al. 2010)

- 6.4.4 Based on visiting rates to each access point and surrounding residential densities, Clarke et al. (2010) constructed a model that could predict the number of additional visitors to the SPA that would result from new residential developments close to the Forest. The predictive model allows for distance to the SPA access points and the characteristics of the access point. Table 6.1 illustrates the model outputs in relation to developments of 100 additional dwellings at a range of settlements around the SPA.
- 6.4.5 The model provides a means to directly compare the consequences of development (in terms of increased SPA visitor numbers) at a potential development location. Accordingly, 100 new dwellings at Crowborough, in close proximity to parts of the SPA, is predicted to lead to 12.2 extra visitors per 16 hours, in contrast to 4.1 extra visitors for an equivalent number of dwellings at East Grinstead, or 1.2 extra visitors for the same number at Haywards Heath, further away

from the Forest (Clarke *et al*, 2010). The analysis shows that, although the existing numbers of visitors to Ashdown Forest may not be negatively affecting populations of Dartford warbler and nightjar, the visitors associated with new strategic housing allocations may do, especially in combination with the effects of other plans and projects. Generally speaking, the closer an individual dwelling or residential development is to the Forest, the more likely its inhabitants are to visit on a regular basis.

Settlement	Distance from SPA *	Number of added visits per 100 dwellings **
East Grinstead	5.10	4.1
Crawley	12.98	0.3
Haywards Heath	9.48	1.2
Uckfield	4.99	3.9
Crowborough	1.50	12.2
Royal Tunbridge Wells	10.25	0.8

Table 6.1: Predicted additional visitor rates to the SPA per 16 daylight hours in Sept.resulting from new development at different locations (Source: Clarke et al. 2010)

* Shortest distance from settlement boundary to SPA boundary – except Crowborough ** Visits per 16 daylight hours in September

- 6.4.6 Clarke *et al* (2010) found that the majority of visitors travelling by car (>85%) originated from within a 15km distance from the Forest (Figure 6.1). In order to establish a zone of influence around Ashdown Forest, the distances between post code origin and SAC/SPA from the 2008 field survey data were recalculated for all modes of transport; see Figure 6.2. The recalculations excluded invalid post codes, stem post codes, records that gave no response and those within the Forest (*n*=286 out of 639 interviews conducted).
- 6.4.7 Following consultations with Natural England, a 7km zone of influence around Ashdown Forest was established; see Figure 6.3. This is the area within which the majority (83%) of regular visitors to the Forest originate, and therefore where measures targeted at reducing pressure on the Forest would be most effective.
- 6.4.8 Natural England has stated that 8ha of Suitable Alternative Natural Greenspace (SANG) should be provided for every 1,000 increase in population (or part thereof) within this zone, in line with the Thames Basin Heaths approach to avoidance and mitigation. SANGs should be complimented by development-funded changes to access management and monitoring within Ashdown Forest to reduce the onsite impacts of the remaining people who will inevitably continue to visit the site. These measures are collectively referred to as Strategic Access Management and Monitoring (SAMM) measures.

6.5 Contribution of the District Plan to Future Visiting Rates within Ashdown Forest

6.5.1 The Consultation Draft District Plan identifies an estimate of objectively assessed housing needs in Mid Sussex. This number is currently 516 per annum (8,776 to 2031) although the Plan



requirement number is still subject to confirmation. This requirement will be delivered through strategic developments at Burgess Hill (outside of the 7km zone of influence) and Neighbourhood Plans. The majority of towns and parishes in the District are committed to preparing their own Neighbourhood Plan, but it is not currently known how many homes could be delivered within the 7km zone around the Ashdown Forest over the plan period.





Figure 6.2: Distances travelled to Ashdown Forest in 2008 by all respondents who gave a full post code (n=286)

6.5.2 Once the likely numbers of homes to be delivered within the 7km zone of influence has been established, it will be possible to calculate the amount of SANG that would be required to support this level of development in order to avoid adverse disturbance effects on the Ashdown Forest SPA's bird assemblage. This can be calculated in accordance with Natural England's requirement for 8ha of SANG for every 1,000 increase in population and an average dwelling occupancy rate of 2.38 (calculated by dividing the district's total resident population in March 2011⁴ (139,860) by the total number of dwellings⁵ (58,712).

⁵ Office for National Statistics: Neighbourhood Statistics [accessed September 2014]: http://www.neighbourhood.statistics.gov.uk/dissemination/LeadTableView.do?a=7&b=6275169&c=mid+sussex&d=13&e=7&g=647 4807&i=1001x1003x1004&m=0&r=1&s=1410950738425&enc=1&dsFamilyId=2512



⁴ Office for National Statistics: Neighbourhood Statistics [accessed September 2014]: http://www.neighbourhood.statistics.gov.uk/dissemination/LeadTableView.do?a=7&b=6275169&c=mid+sussex&d=13&e=13&g=64 74807&i=1001x1003x1004&m=0&r=1&s=1410950623831&enc=1&dsFamilyId=2491



6.6 Other Plans and Projects acting In Combination

- 6.6.1 Other plans and projects which may act in combination with the District Plan to exacerbate disturbance impacts to the SPA include proposed residential developments resulting in a net gain in dwellings within the 7km zone of influence in the following authority areas:
 - Lewes District Council;
 - Sevenoaks District Council;
 - > Tandridge District Council;
 - > Tunbridge Wells Borough Council; and
 - Wealden District Council.
- 6.6.2 It is understood that local planning authorities in Lewes, Tunbridge Wells and Wealden have similar arrangements in place regarding the provision of SANG and SAMM measures, and that no strategic housing sites are currently being considered in Sevenoaks and Tandridge districts. In any event, Mid Sussex District Council's responsibility is limited to avoiding and mitigating the impacts of developments proposed within its own area.

6.7 Avoidance and Mitigation Measures proposed by the District Plan

6.7.1 Responding to the findings and recommendations of earlier versions of the HRA for the District Plan, and its forerunner the Core Strategy, the current District Plan seeks to avoid and mitigate disturbance impacts to Ashdown Forest SPA through policy DP14; see Box 2.

Box 2: Policy DP14 – Ashdown Forest Special Area of Conservation and Special Protection Area

In order to prevent adverse effects on the Ashdown Forest SPA and SAC, the Council will work with partners to develop a strategic approach to protecting it from recreational pressures. New residential development will be required to demonstrate that adequate measures are put in place to avoid or mitigate any potential adverse effects. Where bespoke mitigation is provided, these measures will need to be in place before occupation of development and be in perpetuity. The following provisions will apply:

1) Within a 400 metres buffer zone around the Ashdown Forest mitigation measures are unlikely to be capable of protecting the integrity of the SPA and therefore residential development will not be permitted.

2) Within a 7km zone of influence around the Ashdown Forest SPA, residential development leading to a net increase in dwellings will be required to contribute to:

- The provision of Suitable Alternative Natural Greenspace (SANGs) to the minimum level of 8Ha per 1,000 net increase in population or in the form of financial contributions to SANGs elsewhere; and
- The Ashdown Forest Strategic Access Management and Monitoring (SAMM) Strategy.
- 6.7.2 The purpose of SANG is to cater for the daily recreational needs of the residents of new (and existing) developments. By providing attractive, accessible greenspace close to home, SANGs thereby reducing the likelihood of residents travelling to Ashdown Forest to satisfy this need. It is acknowledged, however, that it will neither be possible nor desirable to prevent all new



residents from visiting the SPA on an occasional basis. Over recent years the Council has been working with Parish and Town Councils within 7km of the SPA to understand their future housing needs, and to identify one or more strategic sites that are capable of meeting the SANG requirement described at section 6.5. The Council has published a guidance note on the required standards for SANGs which is available on its website⁶. A press release on the Council's website⁷ announces recent confirmation of an to extension the Council's agreement to manage and maintain Ashplats Wood and East Court on behalf of East Grinstead War Memorial Limited. The agreement, which will run for at least 125 years, will result in Ashplats Wood being designated and managed as SANG. Initial improvements planned for the 40ha site to enhance its capacity to meet SANG standards include a surfaced 2.5 kilometre circular path that will provide a year-round walking surface for local people. Further improvements are planned for the future and the Council and its partners are preparing a rolling 10 year conservation and management plan for the site.

- 6.7.3 The purpose of the Strategic Access Management and Monitoring Strategy is to mitigate the adverse effects of disturbance caused by the proportion of visits from new residential developments close to the SPA which will not be avoided through the provision of SANG. The Council has developed an interim SAMM Strategy⁸ to implement the recommendations of earlier versions of the HRA. This has been in place since August 2013 and sets out mitigation measures which have been discussed and agreed in collaboration with the Conservators of Ashdown Forest, Natural England and the other affected local authorities (Lewes District Council, Wealden District Council and Tunbridge Wells Borough Council). These measures focus on protecting the SPA from new recreational pressures through managing access (visitor) behaviour and monitoring both birds and visitors. The interim SAMM Strategy will be superseded by a joint SAMM Strategy with Wealden District Council, Lewes District Council and Tunbridge Wells Borough Council, and this work is underway.
- 6.7.4 Development contributions are being collected towards the interim SAMM Strategy via Section 106 of the Town and Country Planning Act 1990. Development contributions towards delivering the strategic SANG(s) for Mid Sussex will be collected once work on finalising the SANG(s) is complete.

6.8 Impact Assessment

6.8.1 This section assesses whether the District Plan can be expected to adversely affect the ecological integrity of the Ashdown Forest SPA as a result of disturbance impacts. It does this by considering the sites' conservation objectives (Table 3.2), with reference to the characteristics of the site (Chapter 4) and its qualifying features (Table 3.1), and in the context of available data

⁸ Mid Sussex District Council (2013): Ashdown Forest Special Protection Area (SPA) and Special Area of Conservation (SAC) Strategic Access Management and Monitoring (SAMM) – Interim Mitigation Strategy [accessed September 2014]: http://www.midsussex.gov.uk/media/SAMMInterimMitigationStrategySept13.pdf



⁶ Mid Sussex District Council (2012): Suitable Alternative Natural Greenspace Guidance [accessed September 2014]: http://www.midsussex.gov.uk/media/SANGCriteriaWebsite_Sept12.pdf

⁷ Mid Sussex District Council (2014): Press Release: Ashplats Wood improvements [accessed September 2014]: http://www.midsussex.gov.uk/9517.htm

on disturbance impacts and the District Plan's policy response as outlined above (sections 6.1 to 6.7).

Objective 1: Extent and distribution of the habitats of the qualifying features

6.8.2 Ashdown Forest is classified as an SPA for its breeding populations of Dartford warbler and nightjar, which are listed on Annex 1 of the Birds Directive. Breeding woodlark (also an Annex 1 species) are present in qualifying numbers. The strategic SANG(s) for Mid Sussex will be designed to avoid the majority of additional recreational visits to the SPA resulting from planned residential developments within the District Plan. Together with the (current) interim SAMM Strategy and (future) joint SAMM Strategy, this will ensure that a sufficient extent and distribution of suitable breeding and foraging habitat (in the absence of excess disturbance) for Dartford warbler and nightjar (and woodlark) is likely to be maintained on a long-term basis.

Objective 2: Structure and function of the habitats of the qualifying features

6.8.3 The commitment towards delivering SANG and the accompanying SAMM Strategy is considered to demonstrate that sufficient measures are available to ensure that there will be no adverse effect on the structure and function of the habitats of qualifying species as a result of increased disturbance impacts.

Objective 3: Supporting processes on which the habitats of the qualifying features rely

6.8.4 The commitment towards delivering SANG and the accompanying SAMM Strategy is considered to demonstrate that sufficient measures are available to ensure that there will be no adverse effect on the supporting processes on which the habitats of qualifying species rely as a result of increased disturbance impacts.

Objective 4: Populations of the qualifying features

6.8.5 The commitment towards delivering SANG and the accompanying SAMM Strategy is considered to demonstrate that sufficient measures are available to ensure that there will be no adverse effect on the populations of qualifying species as a result of increased disturbance impacts.

Objective 5: Distribution of qualifying features within the site

6.8.6 The commitment towards delivering SANG and the accompanying SAMM Strategy is considered to demonstrate that sufficient measures are available to ensure that there will be no adverse effect on the distributions of qualifying species within the site as a result of increased disturbance impacts.

6.9 Conclusions regarding Adverse Effects on Integrity

6.9.1 It can be concluded that the District Plan will not result in adverse effects on the ecological integrity of the Ashdown Forest SPA.



This page is intentionally blank.

7 Summary and Conclusions

7.1 Summary

- 7.1.1 This document sets out the Habitats Regulations Assessment for the Consultation Draft District Plan for Mid Sussex. It draws on information previously published in other documents which form part of the overall HRA procedure for the District Plan, including research reports and earlier versions of the HRA.
- 7.1.2 The assessment addresses the following proposed policies within the District Plan:
 - DP2 Sustainable Economic Development;
 - DP3 Town Centre Development;
 - DP5 Housing;
 - DP7 Strategic Allocation to the east of Burgess Hill at Kings Way;
 - > DP8 Strategic Allocation to the north and north-west of Burgess Hill;
 - DP12 New Homes in the Countryside;
 - DP29 Rural Exception Sites; and
 - DP30 Gypsies, Travellers and Travelling Showpeople.

7.2 Conclusions

- 7.2.1 It can be concluded that the District Plan will not result in adverse effects on the ecological integrity of the Ashdown Forest SAC.
- 7.2.2 It can be concluded that the District Plan will not result in adverse effects on the ecological integrity of the Ashdown Forest SPA.

7.3 Consultation Arrangements

7.3.1 The Consultation Draft District Plan, along with the HRA Report, will be made available for public consultation. If you wish to comment on the documents, your comments should be submitted using the contact details overleaf.

7.4 Next Steps

7.4.1 Following consultation on the Consultation Draft District Plan and its Habitats Regulations Assessment, the Plan will move towards Publication and Submission. Representations received on the Draft Plan may result in changes to the plan and, if so, significant changes will also be



subject to further appraisal. This may lead to a further edition of the HRA Report being produced, or perhaps an Addendum to the current report, so that the findings of further appraisal can be circulated. The representations, proposed changes to the plan and HRA information will be submitted to the Planning Inspector to consider during the Examination in Public.

Email:	
	LDFconsultation@midsussex.gov.uk
Online:	
	www.midsussex.gov.uk/districtplan
By post:	
	Planning Policy and Economic Development Mid Sussex District Council Oaklands Road Haywards Heath West Sussex RH16 1SS



References and Bibliography

Amey (2012): Mid Sussex Transport Study Report.

Baillie, S.R., Marchant, J.H., Leech, D.I., Renwick, A.R., Eglington, S.M., Joys, A.C., Noble, D.G., Barimore, C., Conway, G.J., Downie, I.S., Risely, K. & Robinson, R.A. (2012). BirdTrends 2011. BTO Research Report No. 609. BTO, Thetford. http://www.bto.org/birdtrends

BirdLife International 2012. *Sylvia undata*. In: IUCN 2012. IUCN Red List of Threatened Species. Version 2012.2. <www.iucnredlist.org>. Downloaded on 6 November 2012.

Bobbink R, Boxman D, Fremstad E, Heil G, Houdijk A and Roelofs J (1993): Nitrogen eutrophication and critical load for nitrogen based upon changes in flora and fauna in (semi)-natural terrestrial ecosystems. In: *Critical loads for nitrogen*. Proceedings of a UN-ECE workshop at Lökeberg, Sweden.

Clarke RT, Sharp J & Liley D (2010): Ashdown Forest Visitor Survey Data Analysis. Natural England Commissioned Reports, Number 048.

Conservators of Ashdown Forest (2013): Strategic Plan for Ashdown Forest (2013-2016).

Conway G, Wotton S, Henderson I, Langston R, Drewitt A & Currie F (2007): Status and distribution of European Nightjars *Caprimulgus europaeus* in the UK in 2004. *Bird Study* **54**, 98–111.

Conway, G., Wotton, S., Henderson, I., Eaton, M., Drewitt, A. & Spencer, J. (2009): The status of breeding Woodlarks *Lullula arborea* in Britain in 2006. *Bird Study* **56**: 310–325.

Department for Communities and Local Government (DCLG, 2006): Planning for the Protection of European Sites: Appropriate Assessment (Draft).

DCLG (2012): National Planning Policy Framework.

Department for the Environment, Food and Rural Affairs (Defra) (March 2012): Report of the Habitats and Wild Birds Directives Implementation Review.

Dodd AM, Cleary BE, Dawkins JS, Byron HJ, Palframan LJ & Williams GM (2007): The Appropriate Assessment of Spatial Plans in England: a guide to why, when and how to do it.

Dore CJ *et al* (2003): *UK Emissions of Air Pollutants* 1970 – 2003. UK National Atmospheric Emissions Inventory.

Eaton MA, Brown AF, Noble DG, Musgrove AJ, Hearn R, Aebischer NJ, Gibbons DW, Evans A and Gregory RD (2009): Birds of Conservation Concern 3: the population status of birds in the United Kingdom, Channel Islands and the Isle of Man. *British Birds*, **102**, pp.296–341.

English Nature (1997a&b, 1999 and 2001): Habitats Regulations Guidance Notes 1 – 4.



English Nature (2004): Internal Guidance to decisions on 'site integrity': A framework for provision of advice to competent authorities.

Environment Agency (2005): Further Guidance on Applying the Habitats Regulations to Integrated Pollution Control (IPC), Pollution Prevention and Control (PPC) and Control of Major Accident Hazards (COMAH), Comprising of Appendix 7A for IPC and PPC and Appendices 7B and 7C for COMAH. Number 37_02.

Environment Agency (2010): *Horizontal Guidance Note H1 – Annex (f)*. Published at <u>www.environment-agency.gov.uk</u>

European Commission (2000a): Communication from the Commission on the Precautionary Principle.

European Commission (2000b): Managing Natura 2000 Sites: The provisions of Article 6 of the Habitats Directive 92/43/EEC.

European Commission (2001): Assessment of plans and projects significantly affecting Natura 2000 Sites: Methodological Guidance on the Provisions of Article 6(3) and 6(4) of the Habitats Directive.

European Commission (2007): Interpretation Manual of European Union Habitats.

European Council (1992): Council Directive 92/43/EEC on the conservation of natural habitats and of wild fauna and flora.

European Council (2009): Council Directive 2009/147/EC on the conservation of wild birds.

Gibbons, D.W., Reid, J.B. & Chapman, R.A. (1993): The New Atlas of Breeding Birds in Britain and Ireland: 1988–1991. T. & A.D. Poyser, London.

Gibbons DW & Wotton S (1996): The Dartford Warbler in the United Kingdom in 1994. *British Birds*, **89**, 203-212.

Hall, R.J.; Emmett, B.; Garbutt, A.; Jones, L.; Rowe, E.; Sheppard, L.; Vanguelova, E.; Pitman, R.; Britton, A.; Hester, A.; Ashmore, M.; Power, S.; Caporn, S. (2011): *UK Status Report July 2011: Update to empirical critical loads of nitrogen*. Report to Defra under contract AQ801 Critical Loads and Dynamic Modelling, cited on http://www.apis.ac.uk/node/972

Highways Agency (2007): Design Manual for Roads and Bridges: Volume 11 Environmental Assessment, Section 3 Environmental Assessment Techniques, Part 1 Air Quality (HA207/07).

James P [ed.] (1996): Birds of Sussex.

Joint Nature Conservancy Council website [accessed September 2014]: <u>http://jncc.defra.gov.uk/ProtectedSites/SACselection/SAC habitats.asp</u>

Langston RHW, Liley D, Murison G, Woodfield E & Clarke RT (2007): What effects do walkers and dogs have on the distribution and productivity of breeding European nightjar Caprimulgus europaeus? Ibis **149** (Suppl. 1): 27 – 36.



Laxen D and Wilson P (2002): A New Approach to Deriving NO_2 from NO_X for Air Quality Assessment of Roads. Report prepared on behalf of Defra and the devolved administrations.

Liley D & Clarke RT (2003): The impact of urban development and human disturbance on the numbers of nightjar *Caprimulgus europaeus* on heathlands in Dorset, England. *Biological Conservation* **114**: 219 – 230.

Liley D and Sutherland WJ (2007): Predicting the population consequences of human disturbance for Ringed Plovers *Charadrius hiaticula*: a game theory approach. *Ibis*, **149**, pp.82-94.

Mallord JW (2005): Predicting the consequences of human disturbance, urbanisation and fragmentation for a woodlark *Lullula arborea* population. *School of Biological Sciences*. UEA, Norwich.

Mallord, J.W., Dolman, P.M., Brown, A.F. & Sutherland, W.J. (2007a): Linking recreational disturbance to population size in a ground-nesting passerine. *Journal of Applied Ecology* **44**: 185–195.

Mallord JW, Dolman PM, Brown AF and Sutherland WJ (2007b): Quantifying density dependence in a bird population using human disturbance. *Oecologia*, **153**, pp.49-56.

Morris A, Burges D, Fuller RJ, Evans AD & Smith KW (1994): The status and distribution of nightjars *Caprimulgus europaeus* in Britain in 1992. A report to the British Trust for Ornithology. *Bird Study*, **41**, 181-191.

Murison G (2002): The Impact of Human Disturbance on the Breeding Success of Nightjar Caprimulgus europaeus on Heathlands in South Dorset, England. English Nature Research Reports No. 483.

Murison G, Bullock JM, Underhill-Day J, Langston R, Brown AF & Sutherland WJ (2007): Habitat type determines the effects of disturbance on the breeding productivity of the Dartford Warbler *Sylvia undata*. *Ibis* **149** (Suppl. 1): 16 – 26.

Natural England (2008): Habitats Regulations Assessment Scoping Report for Mid Sussex Core Strategy Pre-Submission Document (Consultation Response).

Natural England (draft, 2008): Guidelines for the creation of Suitable Accessible Natural Green Space. Version 12/06/08.

Nilsson J and Grennfelt P (Eds) (1988):Critical Loads for Sulphur and Nitrogen.Quoted by the AirPollutionInformationSystem,accessedonlineat(14/09/09):http://www.apis.ac.uk/overview/issues/overview/cloadslevels.htm

Office of the Deputy Prime Minister (ODPM) (2005): Government Circular: Biodiversity and Geological Conservation – Statutory Obligations and their Impact within the Planning System.

Pers. comm. (2008a; 23/12/2008): Ashdown Forest Car Park Counter and Bird Trend Data, provided by Hew Prendergast, Superintendent, Conservators of Ashdown Forest.

Pers. comm. (2008b; 15/10/2008): Ashdown Forest SAC Deposition Data, provided by Colin Powlesland, Air Quality and Health Policy Manager, Environment Agency.



Pers. comm. (2009; 01/04/2009): National Bird Trends, provided by Carrie Temple, Senior Conservation Officer, RSPB.

Phoenix, G.K.; Emmett, B.A.; Britton, A.J.; Caporn, S.J.M.; Dise, N.B.; Helliwell, R.; Jones, M.L.M.; Leake, J.R.; Leith, I.D.; Sheppard, L.J.; Sowerby, A.; Pilkington, M.G.; Rowe, E.C.; MR, A.; Power, S.A. (2012): Impacts of atmospheric nitrogen deposition: responses of multiple plant and soil parameters across contrasting ecosystems in long-term field experiments. *Global Change Biology* **18** 1197-1215, cited on http://www.apis.ac.uk/node/972

Pitcairn CER, Fowler D and Grace J (1991): Changes in species composition of semi-natural vegetation associated with the increase in atmospheric inputs of nitrogen. *Report to Nature Conservancy Council*. Institute of Terrestrial Ecology.

Rowe, E.C.; Jones, M.L.M.; Henrys, P.A.; Smart, S.M.; Tipping, E.; Mills, R.T.E.; Evans, C.D. (2011): Predicting effects of N pollutant load on plant species based on a dynamic soil eutrophication indicator. Science Report 39, cited on <u>http://www.apis.ac.uk/node/972</u>

Sitters, H.P., Fuller, R.J., Hoblyn, R.A., Wright, M.T., Cowie, N. & Bowden, C.G.R. (1996): The Woodlark *Lullula arborea* in Britain: population trends, distribution and habitat occupancy. *Bird Study* **43**: 172–187.

Transport and Travel Research Ltd (2005): Best Practice Guide for Assessment of Traffic and Air Quality Impacts. Prepared for The West London Air Alliance Quality Cluster Group, with Bureau Veritas.

Tyldesley D (2009): The Habitats Regulations Assessment of Local Development Documents: Revised Draft Guidance for Natural England.

UE Associates and University of Brighton (2009): Visitor Access Patterns on the Ashdown Forest: Recreational use and nature conservation.

UE Associates (2011): Wealden District (incorporating part of the South Downs National Park) Local Development Framework: Assessment of the Core Strategy under the Habitats Regulations – Submission Stage.

Underhill-Day JC (2005): A literature review of urban effects on lowland heaths and their wildlife. English Nature Research Reports, No. 623.

Underhill-Day JC & Liley D (2007): Visitor patterns on southern heaths: a review of visitor access patterns to heathlands in the UK and the relevance to Annex I bird species. *Ibis* **149** (Suppl. 1): 112 – 119.

United Nations Educational, Scientific and Cultural Organisation (UNESCO, 1971): Convention on Wetlands of International Importance especially as Waterfowl Habitat. (Ramsar (Iran), 2 February 1971, UN Treaty Series No. 14583).

Urban Edge Environmental Consulting (2013): Habitats Regulations Assessment for the Mid Sussex District Plan: May 2013.

West Sussex County Council (2011): West Sussex Transport Plan 2011-2026.



Wotton, S.R. & Gillings, S. (2000): The status of breeding Woodlarks *Lullula arborea* in Britain in 1997. *Bird Study* **47**: 212–224.

Wotton S, Conway G, Eaton M, Henderson I & Grice P (2009): The status of the Dartford Warbler *Sylvia undata* in the UK and the Channel Islands in 2006. *British Birds*. **102**, pp.230-246.

Wright, L.J., Hoblyn, R.A., Sutherland, W.J. & Dolman, P.M. (2007): Reproductive success of Woodlarks *Lullula arborea* in traditional and recently colonized habitats. *Bird Study* **54**: 315–323.

Wright, L.J., Hoblyn, R.A., Green, R.E., Bowden, C.G.R., Mallord, J.W., Sutherland, W.J. & Dolman, P.M. (2009): Importance of climatic and environmental change in the demography of a multi-brooded passerine, the Woodlark *Lullula arborea*. *Journal of Animal Ecology* **78**: 1191–1202.



This page is intentionally blank.

Appendix I: Updated Screening Assessment

The following table presents the updated findings of the HRA screening exercise, which categorises each policy proposal according to the likelihood of it leading to significant effects on a European site. The key which follows the table describes the colours and alphanumeric coding for each category.

No.	Policy title	Ashdown Forest	
		SAC	SPA
DP1	Sustainable development in Mid Sussex	A1	A1
DP2	Sustainable economic development	D2	A4
DP3	Town centre development	D2	A4
DP4	Village and Neighbourhood Centre development	В	В
DP5	Housing	D2	D2
DP6	General principles for strategic development at Burgess Hill	A1	A1
DP7	Strategic allocation to the east of Burgess Hill at Kings Way	D2	A4
DP8	Strategic allocation to the north and northwest of Burgess Hill	D2	A4
DP9	Protection and enhancement of countryside	A1	A1
DP10	Preventing coalescence	A1	A1
DP11	Sustainable rural development and the rural economy	A1	A1
DP12	New homes in the countryside	A1	D2
DP13	High Weald Area of Outstanding Natural Beauty	A3	A3
DP14	Ashdown Forest SAC and SPA	A4	A4
DP15	Setting of the South Downs National Park	A3	A3
DP16	Sustainable tourism	A1	A1
DP17	Securing infrastructure	A1	A1
DP18	Transport	A1	A1
DP19	Rights of Way and other recreational routes	A1	A1
DP20	Communication infrastructure	A1	A1
DP21	Leisure and cultural facilities and activities	A1	A1
DP22	Community facilities and local services	A1	A1
DP23	Character and design	A1	A1
DP24	Dwelling space standards	A1	A1
DP25	Accessibility	A1	A1

No.	Policy title	Ashdown Forest	
		SAC	SPA
DP26	Noise, air and light pollution	A3	A3
DP27	Housing mix	A1	A1
DP28	Affordable housing	A1	A1
DP29	Rural exception sites	A1	D2
DP30	Gypsy, travellers and travelling showpeople	A1	D2
DP31	Listed Buildings and other buildings of merit	A3	A3
DP32	Conservation Areas	A3	A3
DP33	Historic Parks and Gardens	A3	A3
DP34	Archaeological sites	A3	A3
DP35	Trees, woodland and hedgerows	A3	A3
DP36	Biodiversity	A3	A3
DP37	Green infrastructure	A1	A1
DP38	Sustainable design and construction	A3	A3
DP39	Renewable energy schemes	A3	A3
DP40	Flood risk and drainage	A3	A3
DP41	Water infrastructure and the water environment	A3	A3

Key: Categories for the screening assessment of policies (derived from Tyldesley, 2009)

Categ	ory A: No negative effect
A1	Options / policies that will not themselves lead to development e.g. because they relate to design or other qualitative criteria for development, or they are not a land use planning policy.
A2	Options / policies intended to protect the natural environment, including biodiversity.
A3	Options / policies intended to conserve or enhance the natural, built or historic environment, where enhancement measures will not be likely to have any negative effect on a European Site.
A4	Options / policies that positively steer development away from European sites and associated sensitive areas.
A5	Options / policies that would have no effect because no development could occur through the policy itself, the development being implemented through later policies in the same plan, which are more specific and therefore more appropriate to assess for their effects on European Sites and associated sensitive areas.
Categ	ory B: No significant effect
В	Options / policies that could have an effect, but the likelihood is there would be no significant negative effect on a European site either alone or in combination with other elements of the same plan, or other plans or projects.

Category C: Likely significant effect alone



C1	The option, policy or proposal could directly affect a European site because it provides for, or steers, a quantity or type of development onto a European site, or adjacent to it.
C2	The option, policy or proposal could indirectly affect a European site e.g. because it provides for, or steers, a quantity or type of development that may be very close to it, or ecologically, hydrologically or physically connected to it or it may increase disturbance as a result of increased recreational pressures.
C3	Proposals for a magnitude of development that, no matter where it was located, the development would be likely to have a significant effect on a European site.
C4	An option, or policy that makes provision for a quantity / type of development (and may indicate one or more broad locations e.g. a particular part of the plan area), but the effects are uncertain because the detailed location of the development is to be selected following consideration of options in a later, more specific plan. The consideration of options in the later plan will assess potential effects on European Sites, but because the development could possibly affect a European site a significant effect cannot be ruled out on the basis of objective information.
C5	Options, policies or proposals for developments or infrastructure projects that could block options or alternatives for the provision of other development or projects in the future, which will be required in the public interest, that may lead to adverse effects on European sites, which would otherwise be avoided.
C6	Options, policies or proposals which depend on how the policies etc are implemented in due course, for example, through the development management process. There is a theoretical possibility that if implemented in one or more particular ways, the proposal could possibly have a significant effect on a European site.
C7	Any other options, policies or proposals that would be vulnerable to failure under the Habitats Regulations at project assessment stage; to include them in the plan would be regarded by the EC as 'faulty planning'.
C8	Any other proposal that may have an adverse effect on a European site, which might try to pass the tests of the Habitats Regulations at project assessment stage by arguing that the plan provides the imperative reasons of overriding public interest to justify its consent despite a negative assessment.
Categ	ory D: Likely significant effects in combination
D1	The option, policy or proposal alone would not be likely to have significant effects but if its effects are combined with the effects of other policies or proposals provided for or coordinated by the LDD (internally) the cumulative effects would be likely to be significant.
D2	Options, policies or proposals that alone would not be likely to have significant effects but if their effects are combined with the effects of other plans or projects, and possibly the effects of other developments provided for in the LDD as well, the combined effects would be likely to be significant.
D3	Options or proposals that are, or could be, part of a programme or sequence of development delivered over a period, where the implementation of the early stages would not have a significant effect on European sites, but which would dictate the nature, scale, duration, location, timing of the whole project, the later stages of which could have an adverse effect on

such sites.

This page is intentionally blank.



Urban Edge Environmental Consulting Ltd

Unit 5 | Westergate Business Centre | Brighton | BN2 4QN T: 01273 68 67 66 | E: enquiries@ueec.co.uk www.ueec.co.uk | 🥑 @UrbanEdgeEnviro © Urban Edge Environmental Consulting Ltd 2014

Urban Edge Environmental Consulting Ltd

Unit 5 | Westergate Business Centre | Brighton | BN2 4QN

T: 01273 68 67 66 | E: enquiries@ueec.co.uk

www.ueec.co.uk | 💓 @UrbanEdgeEnviro

© Urban Edge Environmental Consulting Ltd 2014

URBAN EDGE ENVIRONMENTAL CONSULTING

NATURAL PROGRESSION